

WISCONSIN DEPARTMENT OF NATURAL RESOURCES
Root River Steelhead Facility
Fall 2021 and Spring 2022
August 2025




Photo credit: Wisconsin DNR

WRITTEN BY
LAURA SCHMIDT
DNR Fisheries Management
Southern Lake Michigan Fisheries Work Unit
600 E. Greenfield Ave.
Milwaukee, WI 53204

Summary – A total of 2,174 Chinook salmon, 2,340 coho salmon, 1,693 steelhead and 12 brown trout were examined during fall 2021 and spring 2022 at the Root River Steelhead Facility (RRSF). A total of 608 coho were spawned, and approximately 325,000 coho eggs were collected for our hatcheries.

The fall 2021 return of 2,174 Chinook salmon fell just above the ten-year average return. The fall 2021 return of 2,340 coho salmon was 37% above the ten-year average. The fall 2021 return of 55 steelhead was the highest return seen since 2010. Twelve brown trout were encountered in the facility in the fall, which is typically shut down for the season before the majority of brown trout are in the Root River.

The spring 2022 return of steelhead was a total of 1,638. This return was the highest on record since 2000 and 48% above the ten-year average. A total of 1,314 steelhead were spawned, and a total of 2,841,100 eggs were collected.

The number of fish captured at RRSF is a subset of the seasonal migration in the Root River. RRSF does not stop every fish in the river, as they are able to move upstream past the facility before it is operational in early spring and fall, and some fish are able to bypass the facility during the sampling season when the river is at high flows. In 2021, high flows on the Root River for much of the early spring as a result of snow melt and heavy rain delayed start up of the facility, and it is probable that steelhead were moving upstream before the facility was running.

In the fall of 2021 at RRSF the standard weight of a 30-inch Chinook salmon was 9.2 pounds, which was slightly lower than the previous year, but about the same as the ten-year average (Figure 1). The standard weight for a 22-inch coho salmon, which has remained mostly stable over the past ten years, was 3.6 pounds. The standard weight for a 22-inch steelhead was 3.7 pounds, which was slightly lower than the previous year but about the same as the ten-year average. The standard weight for a 20-inch brown trout was 3.5 pounds.

The following tables and figures report the results of data collected at the RRSF during the fall of 2021 and spring 2022. These data contribute to a long-term index of Chinook, coho and steelhead populations in the Root River and are collected to fulfill three objectives: 1) track the abundance of salmonid returns, 2) measure the growth and condition of each species and/or strain, and 3) estimate return rate of each species. For a complete description of methods and calculations, see Thompson and Eggold (2007).

REFERENCES

Thompson, J. and B. Eggold. 2007. Root River Steelhead Facility, Fall 2006 and Spring 2007. Publication number PUB-FH-836 2007. Wisconsin Department of Natural Resources, Milwaukee, Wisconsin. 22 pages.

Table 1. Summary of Chinook salmon, coho salmon, steelhead and brown trout captured at the Root River Steelhead Facility during 2012 to 2022.

HARVEST YEAR	HARVESTED	PASSED UPSTREAM	MISC. SAMPLES	TOTAL
Chinook Salmon				
Fall 2012	201	1,462	167	1,830
Fall 2013	486	1,070	392	1,948
Fall 2014	533	1,646	44	2,223
Fall 2015	384	880	45	1,309
Fall 2016	518	1,375	152	2,045
Fall 2017	696	1,542	524	2,762
Fall 2018	794	326	5	1,125
Fall 2019	578	818	36	1,432
Fall 2020	727	2,889	311	3,927
Fall 2021	417	1,635	122	2,174
Coho Salmon				
Fall 2012	153	659	37	849
Fall 2013	216	1,281	169	1,666
Fall 2014	60	1,295	33	1,388
Fall 2015	60	1,351	25	1,436
Fall 2016	60	1,324	133	1,517
Fall 2017	66	2,290	180	2,536
Fall 2018	61	2,397	29	2,487
Fall 2019	60	1,117	38	1,215
Fall 2020	30	1,508	14	1,552
Fall 2021	39	2,263	38	2,340
Steelhead				
Fall 2012	0	16	2	18
Spring 2013	120	179	2	301
Fall 2013	0	7	0	7
Spring 2014	120	852	7	979
Fall 2014	0	11	0	11
Spring 2015	60	711	3	774
Fall 2015	0	9	0	9
Spring 2016	60	1,293	0	1,353
Fall 2016	0	43	0	43
Spring 2017	120	774	1	895
Fall 2017	0	9	0	9
Spring 2018	62	990	0	1,052
Fall 2018	0	20	0	20
Spring 2019	91	651	0	742
Fall 2019	8	13	0	21
Spring 2020	18	110	0	128
Fall 2020	8	2	0	10
Spring 2021	167	429	0	596
Fall 2021	40	15	0	55
Spring 2022	222	1,408	8	1,638
Brown Trout				
Fall 2012	0	112	22	134
Fall 2013	0	166	1	167
Fall 2014	13	174	5	192
Fall 2015	0	52	34	86
Fall 2016	0	11	5	16
Fall 2017	0	12	3	15
Fall 2018	0	80	0	80
Fall 2019	0	9	0	9
Fall 2020	0	2	0	2
Fall 2021	0	12	0	12




Figure 1. Standard weight for the major salmonid species returning to the Root River Steelhead Facility from 2012 to 2021. For brown trout, breaks in the graph represent years where returns were too low to estimate standard weight.

CHINOOK SALMON

In fall 2021, 75% of Chinook salmon handled were passed upstream of the dam after processing (Table 1). Of the fish not passed upstream, the majority were kept for coded-wire tag collection.

Analysis of length-weight data (Table 4) revealed that average length (32.4 inches) and weight (13.1 pounds) of returning Chinook salmon decreased from the previous year but were within the ten-year average. The standard weight (9.2 pounds) was only slightly lower than the previous fall (9.6 pounds) and was about the same as the ten-year average.

Throughout the fall season, Chinook salmon were sub-sampled as part of two ongoing studies. From 2011-2017, the U.S. Fish and Wildlife Service's mass-marking program marked all Chinook salmon stocked into Lake Michigan with an adipose clip and coded-wire tag (CWT) for analysis of movement patterns of Chinooks in the lake, growth rates, natural reproduction, and "straying", when a mature fish returns to a stream other than the one where it was originally stocked.

In addition, from 2015-2018, the Wisconsin DNR conducted a study to evaluate net pens, a collaborative project where fishing clubs hold Chinook salmon in net pens to acclimate them to the rivers. Chinook salmon in the Kewaunee and Root Rivers were differentially marked with coded-wire tags. Chinook stocked directly into the rivers and into net pens received different CWT numbers, and analysis of these tags will help evaluate whether or not Wisconsin's collaborative net pen projects are having a positive impact on post stocking survival.

Tags were recovered from 331 coded-wire tagged Chinooks at RRSF in fall 2021.

From 2012-2021, a total of 4,809 tags were recovered from Chinook salmon, which provides insight into the age structure of Chinook returning to the river. On average, 75% of tagged Chinook that were sampled were ages 2 or 3 (Table 2). In some years, a high number of age-1 Chinooks returned (2016, 2017). In 2021, no age 1 Chinooks were sampled because there were no tagged age-1 fish, and 97% of the tags recovered were from Chinooks ages 2 and 3.

Table 2. Age distribution of coded-wire tagged Chinook salmon (sexes combined) examined at the Root River Steelhead Facility during fall, 2012 through 2021.

YEAR OF RETURN	NUMBER AT AGE							TOTAL NUMBER
	0	1	2	3	4	5	6	
2012	-	49	134	-	-	-	-	183
2013	1	231	103	125	-	-	-	460
2014	4	20	285	160	7	-	-	476
2015	10	31	52	278	1	-	-	372
2016	6	254	68	125	24	-	-	477
2017	-	352	239	81	3	-	-	675
2018	-	144	361	213	3	-	-	721
2019	-	72	276	162	2	1	-	513
2020	-		241	353	6	-	1	601
2021	-	-	1	319	11	-	-	331

Coded-wire tag data also revealed a large overlap in length-at-age for tagged Chinook salmon returning to the weir (Figure 2, Figure 3). In some years (2012, 2015, and 2019), age-1 Chinooks were on average 5-8 inches smaller than older fish, which could possibly be explained by alewife year-class strength. However, in most years, length could not be used as an indicator of age.




Figure 2. Length-at-age of coded-wire tagged Chinook salmon recovered at RRSF from 2012-2017.




Figure 3. Length-at age of coded-wire tagged Chinook salmon recovered at RRSF from 2018-2021.

Coded-wire tag returns also show relatively low rates of “straying” by Chinook salmon into the Root River during the fall spawning run. Not all Chinooks stocked into the Root River by the Wisconsin DNR from 2012-2018 received a unique CWT lot code, primarily due to constraints in hatchery space. In Figures 4 and 5, the “WI South” location includes the Root River, Racine Harbor, Milwaukee Harbor, Port Washington, Sauk Creek, and the Sheboygan River. The “WI North” location includes the Manitowoc River, the East Twin River, and the West Twin River. From 2012-2015, over 40% of tagged Chinooks that were recovered were stocked into the Root River (Figure 4). Almost 20% were stocked in the Pike River. The Pike River can become sandblocked, in which case these fish return to other streams, such as the Root River.

The Root River, Pike River, and Wisconsin south locations combined comprise 88.4% of Chinook returns.




Figure 4. Stocking locations of tagged Chinooks sampled at RRSF from 2012-2015.

From 2016-2021, the Root River, Pike River and Wisconsin south locations combined comprised 97% of sampled Chinook returns (Figure 5). This includes Chinook that were stocked directly into the Root River, and those that were stocked into net pens.




Figure 5. Stocking locations of tagged Chinooks returning to the Root River weir from 2016-2021.

Overall, from 2016-2021, the results of the DNR's net pen study were mixed (Figure 6). The figure below shows the recovery rate (number of fish recovered per number stocked) of Chinook stocked into the Root River or into net pens by year-class. Chinook salmon were tagged in 2015, but first returned to the Root River in 2016. The 2015 and 2016 year-classes stocked directly into the Root River returned at a higher rate than those stocked in net pens. For the 2017 year-class, the return rate was basically the same. For the 2018 year-class, fish stocked directly into net pens returned at a slightly higher rate until fall 2021, when 319 fish directly stocked into the Root River were recovered, but only 3 that were stocked into net pens.




Figure 6. Comparison of tagged Chinooks recovered at RRSF that were direct stocked into the Root River (blue bars) and stocked into net pens (orange bars).

COHO SALMON

In fall 2021, 97% of coho salmon handled were passed upstream of the dam after processing (Table 1).

Analysis of length-weight data (Table 4) revealed that average length (22.1 inches) and average weight (4.3 pounds) of returning coho decreased from the previous year, although the standard weight (3.6 pounds) was slightly higher than the previous fall (3.5 pounds) and was the same as the ten-year average.

The age composition (based on length-frequencies) indicated that the 2021 run was comprised of 36% age 1+ and 64% age 2+ coho salmon (Table 3).

Table 3. Estimated age composition of coho salmon (sexes combined) examined at the Root River Steelhead Facility during fall, 2011 through 2021. Ages were assigned by length-frequency of measured fish.

YEAR OF RETURN	PERCENT AGE COMPOSITION		NUMBER USED IN ANALYSIS	TOTAL RETURN
	1+	2+		
2011	6%	94%	761	1,628
2012	21%	79%	715	849
2013	5%	95%	786	1,666
2014	6%	94%	1,353	1,388
2015	9%	91%	1,161	1,436
2016	29%	71%	1,042	1,517
2017	5%	95%	1,249	2,536
2018	5%	95%	1,746	2,487
2019	15%	85%	1,178	1,215
2020	10%	90%	1,061	1,551
2021	36%	64%	1,841	2,340

STEELHEAD

In fall 2021, 15 of the 55 steelhead captured were passed upstream of the dam after processing (Table 1). The remainder were kept for coded-wire tag analysis, described below.

All steelhead that are stocked into the broodstock rivers (the Root and Kewaunee Rivers) are marked with differential fin clips to identify strain for purposes of spawning. Prior to 2017, each strain (Chambers Creek and Ganaraska) were given fin clips on a 3-year rotational basis to assist in estimating age composition.

In 2018, after completion of the Chinook salmon study, the U.S. Fish and Wildlife Service's mass-marking program began marking all rainbow trout stocked into Lake Michigan with an adipose-clip and coded-wire tags, which will provide data for analysis of movement, growth rates, and estimates of natural reproduction.

To simplify the mass marking process, in which all steelhead are adipose-clipped through the USFWS mass marking trailer, but broodstock fish are hand-clipped for strain identification, a standard fin clip per strain was chosen. As the years of data collection continue, a length-age key will be developed to estimate ages of returned fish to the weir.

Skamania strain steelhead, a fall-run strain, were stocked in 2018 and 2019 (2017 and 2018 year-classes) after obtaining eggs from Indiana. These fish were given an adipose-clip and right maxillary-clip for identification in the field.

Of the 55 steelhead processed in fall 2021, 29 (53%) were Skamania strain, 10 (18%) were Arlee strain, 12 (22%) were Chambers Creek strain, and the remaining 4 (7%) were unknown strain. The 12 Chambers Creek fish were identified as stocked in the Root River by their differential fin clip (adipose + left maxillary) and were passed upstream of the dam. Two of the unknown strain fish had no fin clips and were presumed to be wild or unclipped strays and were also passed upstream.

The Arlee strain was stocked into Wisconsin waters of Lake Michigan from 2001-2020. Approximately 100,000 were stocked per year, split between the northern and southern ports. In southern Lake Michigan, fish were stocked in either the Root River (Racine) and Port Washington or Sheboygan, Milwaukee and Kenosha on an alternating year basis. Thus, in figure 7 below, "Racine/Port Wash" refers to Arlees stocked in Racine and Port Washington, and "Sheb, Milw, Kenosha" refers to Arlees stocked in Sheboygan, Milwaukee and Kenosha.

Over half (64%) of steelhead that were collected for analysis were stocked into the Root River. An additional 19% were Arlee strain that were stocked in either Racine or

Port Washington. The remaining strays were primarily from Illinois and Wisconsin waters, with a few fish returning from Indiana waters (Figure 7).




Figure 7. Stocking locations of CWT Steelhead returning to the Root River weir in fall 2021.

Six Skamania were age 4 (2017 year-class), while 20 were age 3 (2018 year-class). Analysis of growth is limited, but 75% of coded-wire tagged steelhead in fall 2021 were age 3, ranging from 22-30 inches. Age 2 steelhead ranged from 21-28 inches, and age 4 fish ranged from 28-34 inches.




Figure 8. Length-at-age of coded-wire tagged steelhead recovered at RRSF in fall 2021.

In spring 2022, 86% of steelhead handled were passed upstream of the dam after processing (Table 1).

Analysis of length-weight data (Table 4) revealed that average length (24.6 inches) and average weight (5.5 pounds) of returning steelhead were higher than the previous year. The standard weight (3.8 pounds) was also higher than the previous spring, and slightly higher than the ten-year average.

Of the 1,638 steelhead processed, 77 (4.7%) were unclipped and presumed to be wild or unclipped strays. Steelhead stocked in 2021 were not clipped due to COVID-19, so many of these could be unclipped age-2 fish. 723 steelhead (44.1%) were Chambers Creek strain marked with an adipose and left maxillary or only a left maxillary clip, identifying them as stocked into the Root River. 519 (31.7%) were Ganaraska strain fish marked with ventral clips, identifying them as stocked into the Root River. The remaining 319 fish (19.5%) were adipose-only, identifying them in the field as strays.

Table 4. Average weight, average length, standard weight (predicted weight at a given length based on a length-weight regression) and trophy weight (95th percentile) for the major salmonid species returning to the Root River Steelhead Facility from fall 2011 to spring 2022. The lengths used for the calculation of standard weight are: 30 inches for Chinook, 22 inches for coho, 22 inches for steelhead and 20 inches for brown trout. Note: brown trout are not listed due to low returns.

SEASON	NUMBER USED IN ANALYSIS	AVERAGE WEIGHT (POUNDS)	AVERAGE LENGTH (INCHES)	STANDARD WEIGHT	TROPHY WEIGHT
CHINOOK SALMON					
2011-12	564	10.9 ± 5.0	30.3 ± 5.0	9.6	18.5
2012-13	694	10.6 ± 3.4	31.5 ± 3.9	8.6	16.0
2013-14	1,085	12.5 ± 6.2	31.6 ± 5.8	9.6	21.5
2014-15	945	11.8 ± 3.0	32.2 ± 3.2	9.0	16.4
2015-16	920	11.7 ± 3.7	31.6 ± 4.2	9.0	16.9
2016-17	870	10.1 ± 5.3	29.5 ± 5.4	9.5	19.1
2017-18	868	11.1 ± 5.3	30.3 ± 5.0	9.5	19.1
2018-19	852	13.8 ± 6.0	32.6 ± 5.6	9.8	22.5
2019-20	1,415	15.3 ± 5.6	33.6 ± 4.5	10.1	24.6
2020-21	1,739	16.7 ± 5.5	34.9 ± 4.3	9.6	24.9
2021-22	1,521	13.1 ± 5.9	32.4 ± 5.6	9.2	22.0
COHO SALMON					
2011-12	786	7.4 ± 2.3	26.9 ± 3.7	3.6	10.3
2012-13	715	4.0 ± 1.7	22.4 ± 3.8	3.5	6.4
2013-14	786	8.2 ± 2.4	28.1 ± 3.4	3.6	11.3
2014-15	1,353	6.2 ± 1.8	25.9 ± 3.3	3.5	8.6
2015-16	1,161	4.5 ± 1.5	23.3 ± 2.9	3.5	7.0
2016-17	1,042	5.2 ± 2.8	23.8 ± 5.0	3.6	9.4
2017-18	1,249	8.3 ± 2.4	27.7 ± 3.2	3.9	11.6
2018-19	1,742	7.8 ± 2.4	26.9 ± 3.3	3.8	11.2
2019-20	1,177	7.4 ± 3.2	26.5 ± 4.7	3.7	11.6
2020-21	1,061	6.9 ± 2.8	26.2 ± 4.3	3.5	10.9
2021-22	1,841	4.3 ± 2.5	22.1 ± 4.7	3.6	8.3
STEELHEAD					
2011-12	247	5.7 ± 1.8	25.4 ± 2.8	3.7	8.9
2012-13	315	4.5 ± 2.1	23.2 ± 4.3	3.6	7.7
2013-14	605	5.6 ± 2.5	24.6 ± 4.3	3.6	9.6
2014-15	779	4.1 ± 1.9	22.6 ± 3.9	3.3	7.6
2015-16	1,047	4.9 ± 1.9	24.0 ± 2.2	3.8	7.3
2016-17	933	6.6 ± 2.2	25.9 ± 3.5	3.8	9.8
2017-18	1,044	6.9 ± 2.1	26.3 ± 2.9	3.9	10.8
2018-19	747	6.7 ± 2.5	25.7 ± 3.4	3.8	11.1
2019-20	143	6.2 ± 2.8	24.9 ± 4.6	4.1	10.3
2020-21	596	4.2 ± 2.5	22.0 ± 4.2	3.8	8.8
2021-22	1,685	5.5 ± 2.1	24.6 ± 3.3	3.7	9.4

APPENDIX A. ROOT RIVER STOCKING NUMBERS

Table A-1. Number of fingerling Chinook salmon stocked in the Root River during 2012-2021. Chinook salmon were marked with an adipose clip and coded-wire tag from 2011 through 2018, and with an adipose clip in 2019. Fish were not marked in 2020 due to the COVID-19 pandemic. Marking with an adipose clip resumed in 2021.

YEAR STOCKED	TOTAL NUMBER	STRAIN	FINCLIP
2012	112,616	Lake Michigan	A-CWT
2013	75,046	Lake Michigan	A-CWT
2014	76,933	Lake Michigan	A-CWT
2015	52,120	Lake Michigan	A-CWT (regular stocking)
	25,640	Lake Michigan	A-CWT (net pen stocking)
2016	50,918	Lake Michigan	A-CWT (regular stocking)
	25,352	Lake Michigan	A-CWT (net pen stocking)
2017	7,467	Lake Michigan	A (regular stocking)
	43,561	Lake Michigan	A-CWT (regular stocking)
	31,300	Lake Michigan	A-CWT (net pen stocking)
2018	51,383	Lake Michigan	A-CWT (regular stocking)
	32,748	Lake Michigan	A-CWT (net pen stocking)
2019	42,626	Lake Michigan	A (regular stocking)
	42,079	Lake Michigan	A (net pen stocking)
2020	101,919	Lake Michigan	None
2021	101,329	Lake Michigan	A

Table A-2. Number of coho salmon stocked in the Root River during 2012 – 2021.

YEAR STOCKED	TOTAL NUMBER	STRAIN	FINCLIP	AGE
2012	75,153	Lake Michigan	None	Spring yearling 1+
	10,968	Lake Michigan	A-CWT	Spring yearling 1+
2013	83,608	Lake Michigan	None	Spring yearling 1+
2014	79,080	Lake Michigan	None	Spring yearling 1+
2015	83,015	Lake Michigan	None	Spring yearling 1+
	10,008	Lake Michigan	None	Fall fingerling 0+
2016	60,021	Lake Michigan	None	Spring yearling 1+
	10,010	Lake Michigan	None	Fall fingerling 0+
2017	76,432	Lake Michigan	None	Spring yearling 1+
	13,001	Lake Michigan	None	Fall fingerling 0+
2018	76,241	Lake Michigan	None	Spring yearling 1+
2019	76,609	Lake Michigan	None	Spring yearling 1+
2020	73,702	Lake Michigan	None	Spring yearling 1+
	26,182	Lake Michigan	None	Fall fingerling 0+
2021	101,509	Lake Michigan	None	Spring yearling 1+

Table A-3. Number of yearling steelhead stocked in the Root River during 2012-2021. Steelhead stocked into Lake Michigan in 2021 were not adipose-clipped or coded-wire tagged due to the COVID-19 pandemic.

YEAR STOCKED	TOTAL NUMBER	STRAIN	FINCLIP
2012	26,998	Chambers Creek	LMLV
	27,031	Ganaraska	BV
2013	26,995	Chambers Creek	LM
	27,116	Ganaraska	ALV
2014	27,118	Chambers Creek	ALM
	29,535	Ganaraska	ARV
2015	31,389	Chambers Creek	LMLV
	31,459	Ganaraska	BV
2016	27,134	Chambers Creek	LM
	28,218	Ganaraska	ALV
2017	28,085	Chambers Creek	ALM
	27,048	Ganaraska	ARV
2018	30,293	Chambers Creek	ALM
	26,252	Ganaraska	ARV
	34,027	Skamania	ARM
	34,511	Skamania	ARM-CWT
2019	8,503	Chambers Creek	A-CWT
	32,034	Chambers Creek	ALM-CWT
	33,884	Ganaraska	ALV-CWT
2020	32,191	Chambers Creek	ALM-CWT
	34,467	Ganaraska	ALV-CWT
2021	33,207	Chambers Creek	LM
	33,176	Ganaraska	LV

Table A-4. Number of brown trout stocked in the Root River during 2011-2020.

YEAR STOCKED	TOTAL NUMBER	STRAIN	FINCLIP
2012	29,695	Seeforellen	ARV
2013	30,561	Seeforellen	ALM
2014	32,100	Seeforellen	ALV
2015	42,743	Seeforellen	ALP
2016	31,690	Seeforellen	ARP
2017	19,122	Seeforellen	A (regular stocking)
	9,383	Seeforellen	A (net pens)
2018	31,448	Seeforellen	A
2019	31,736	Seeforellen	A
2020	32,066	Seeforellen	A
	4,996	Seeforellen	None-fall fingerling
2021	34,074	Seeforellen	A