

WISCONSIN DEPARTMENT OF NATURAL RESOURCES

AMNICON LAKE

2024 – 2025 CREEL SURVEY REPORT


DOUGLAS COUNTY



Treaty Fisheries Publication



Compiled by
Chris Coffin
Fisheries Technician



CONTENTS

INTRODUCTION

General Lake Information	4
Location	4
Physical Characteristics	4
Seasons Surveyed.....	4
Weather	4
Fishing Regulations.....	4
Species Catch And Harvest Information	4
Creel Survey Results And Discussion	5
Survey Logistics.....	5
General Angler Information	5
Results By Species	5
Acknowledgments.....	6

SUMMARY TABLES

Table 1. Sportfishing effort summary	7
Table 2. Creel survey synopses.....	8

SPECIES CATCH AND HARVEST FIGURES

Gamefish	
Figure 1. Walleye	9
Figure 2. Northern Pike.....	10
Figure 3. Muskellunge	11
Figure 4. Smallmouth Bass.....	12
Figure 5. Largemouth Bass	13
Panfish	
Figure 6. Yellow Perch	14
Figure 7. Bluegill.....	15
Figure 8. Black Crappie.....	16
Figure 9. Pumpkinseed	17
Figure 10. Rock Bass.....	18

INTRODUCTION

Fish populations can fluctuate due to a variety of factors including natural forces like climate, reproductive success, predation, and competition. Human activities such as fish harvest, stocking, habitat change, and invasive species introduction can also have significant impacts. Wisconsin Department of Natural Resources (DNR) fisheries crews regularly conduct fishery surveys on lakes and reservoirs to gather the information needed to monitor changes, identify concerns, evaluate past management actions, and to prescribe fishery management strategies. Netting and electrofishing surveys are used to gather data on the status of fish populations and communities, measuring such parameters as species composition, population size, reproductive success, size and age distribution and growth rates.

The other key component of the fishery that we often need to measure is angler harvest to assess its impact on the fishery.

On many lakes in the Ceded Territory of northern Wisconsin, the harvest of fish is divided between sport anglers and the six Ojibwe tribes who harvest fish under rights granted by federal treaties. The tribes harvest fish mostly using a highly efficient method, spearing, during a relatively short time period in the spring. Every fish in the spear harvest is counted – a complete “census” of the harvest.

It would be highly impractical and very costly to conduct a complete census of every angler who fishes on a lake. Therefore, we conduct creel surveys.

A creel survey is an assessment tool used to sample the fishing activities of anglers on a body of water and make projections, or estimates, of harvest and other fishery parameters. Creel survey clerks work on randomly selected days and shifts, forty hours per week. The survey is conducted during the open season for gamefish from the first Saturday in May through the first Sunday

in March. Creel surveys are generally not conducted in November when fishing effort is low, and ice conditions are often unsafe. The survey is run during daylight hours, and shift times change from month to month as day length changes.

Creel survey clerks travel their lakes using a boat or snowmobile to count the number of anglers at predetermined times, and to interview anglers who have completed their fishing trip. Data are collected on what species they fished for, catch, harvest, lengths of fish harvested, marks (fin clips or tags), and hours of fishing effort. Collecting information at the end of a fishing trip provides the most accurate assessment of angling activities, and it avoids the need to disturb anglers while they are fishing.

A computer program is used to estimate catch and harvest of each species, catch and harvest rates, and fishing effort by month, as well as for the year in total. Accurate estimates require that we sample a sufficient and representative portion of the angling activity on a lake. The accuracy of creel survey results depends on good cooperation and truthful responses by anglers when a creel clerk interviews them.


You may have encountered a DNR creel survey clerk on a recent fishing trip. We appreciate your cooperation during an interview. The survey only takes a few minutes of your time, and it gives the DNR valuable information needed for management of the fishery.

This report provides estimates of:

1. Overall fishing effort (pressure)
2. Fishing effort directed at each species
3. Numbers of fish caught and harvested
4. Catch and harvest rates

Also included are a physical description of the lake, discussion of results of the survey, and detailed summaries of fishing effort, catch and harvest.

GENERAL LAKE INFORMATION



LOCATION

Amnicon Lake is located in Douglas County near the town of Superior.

PHYSICAL CHARACTERISTICS

Amnicon Lake is a 426-acre drainage lake with a maximum depth of 31 feet. Littoral substrate consists primarily of muck, with lesser amounts of sand, and gravel. Amnicon Lake contains stained water of moderate transparency.

SEASONS SURVEYED

The open-water creel survey ran from May 4 through Oct. 31, 2024, and the ice fishing creel survey ran from Dec. 1, 2024 through March 2, 2025.

WEATHER

Ice-out on Amnicon Lake was around mid-March, 2024. Fishable ice formed in early December.

FISHING REGULATIONS

The following seasons, daily bag limits, and length limits were in place during the surveyed season:

SPECIES	SEASON	BAG LIMIT	MIN. SIZE
Largemouth Bass	5/ 4-3/ 2	5	14"
Smallmouth Bass	6/ 15-3/ 2	5	14"
Largemouth and Smallmouth Bass Catch and release only all other times of year			
Musky	5/ 25-12/ 31	1	40"
Northern Pike	5/ 4-3/ 2	5	none
Walleye	5/ 4-3/ 2	3	18"
Panfish	year round	25	none

SPECIES CATCH AND HARVEST INFORMATION

Summaries of angling effort, catch and harvest information for each species are in Table 2 and Figures 1-10, along with a comparison of these statistics with the previous creel survey in Table 2, if available. Information about species with fishing seasons extending beyond the season surveyed should be considered minimum estimates. Each species page has up to five graphs depicting the following:

- ESTIMATED FISHING EFFORT**
The estimated number of hours during each month that anglers spent fishing for a species.
- ESTIMATED CATCH AND HARVEST**
The estimated number of fish of the indicated species caught or harvested by all anglers, regardless of targeted species.
- ESTIMATED SPECIFIC CATCH AND HARVEST RATES**
The estimated number of hours it takes an angler to catch or harvest a fish of the indicated species. Only information from anglers who were specifically targeting that species is reported.
- LENGTH DISTRIBUTION OF HARVESTED FISH**
All fish of a species that were measured by the clerk during the entire creel survey season.
- LARGEST AND AVERAGE LENGTH OF HARVESTED FISH**
The largest and average length of a species of fish harvested that month. Only fish measured by the creel survey clerk are reported.

CREEL SURVEY RESULTS AND DISCUSSION

SURVEY LOGISTICS

We encountered no unusual problems conducting the survey or calculating the projections contained in the report.

GENERAL ANGLER INFORMATION

Anglers spent 9,351 hours, or 22 hours per acre, fishing Amnicon Lake during the 2024-2025 season (Table 1). That was less than the Douglas County average of 22.4 hours per acre, and less than the fishing effort documented during the 2006-2007 creel survey (25.9 hours per acre). February was the most heavily fished month (2,194 hours), and fishing effort was lightest in October (266 hours). The creel clerks were able to conduct 434 interviews throughout the survey.

RESULTS BY SPECIES

WALLEYE (Table 2, Figure 1)

Walleye received 6.9 percent of the fishing effort during the season. Anglers spent 928 hours targeting Walleye. The greatest fishing effort for Walleye was in December (174 hours). October had the least amount of Walleye fishing effort (6 hours).

Total catch of Walleye was 40 fish, with a harvest of 15. Highest catch (25 fish) occurred in July, and highest harvest (8 fish) occurred in July. Anglers fished an average of 35.3 hours to catch and 59.5 hours to harvest a Walleye during the survey. The mean length of harvested Walleye was 20.9 inches, and the largest measured was a 26.2-inch fish.

NORTHERN PIKE (Table 2, Figure 2)

Fishing effort directed at Northern Pike was 4,796 hours during the season. Northern Pike fishing effort was greatest in February (1,606 hours). Total catch of Northern Pike was 4,002 fish, with a harvest of 629. Anglers fished an average of 2 hours to catch a Northern Pike during the survey. The mean length of harvested Northern Pike was 22.3 inches, and the largest measured was a 29.9-inch fish.

MUSKELLUNGE (Table 2, Figure 3)

Anglers spent 2,162 hours targeting Muskellunge during the season. Muskellunge fishing effort was greatest in July (629 hours). Total catch of Muskellunge was 83 fish, and the highest catch (26 fish) occurred in July. Anglers fished 27.9 hours to catch a Muskellunge, and there was no documented harvest during the survey.

SMALLMOUTH BASS (Table 2, Figure 4)

Fishing effort targeted at Smallmouth Bass was 17 hours during the season. Smallmouth Bass fishing effort was greatest in August (17 hours). There was no documented catch or harvest of Smallmouth Bass during the survey.

LARGEMOUTH BASS (Table 2, Figure 5)

Fishing effort directed at Largemouth Bass was 1,679 hours during the season. Largemouth Bass fishing effort was greatest in July (558 hours). Total catch of Largemouth Bass was 1,012 fish, with a harvest of 35. Highest catch (391 fish) occurred in July. Anglers fished an average of 2.7 hours to catch a Largemouth Bass during the survey.

PANFISH (Table 2, Figures 6-10)

YELLOW PERCH received 38 hours of directed fishing effort. Total catch of Yellow Perch was 225 fish, with 14 harvested. The mean length of harvested fish was 6 inches.

BLUEGILL received 1,273 hours of directed fishing effort. Total catch of Bluegill was 2,925 fish, with 564 harvested. The mean length of harvested fish was 6.5 inches.

BLACK CRAPPIE received 2,571 hours of directed fishing effort. Anglers caught 5,767 Black Crappie and harvested 777. The mean length of harvested fish was 8 inches.

PUMPKINSEED received no hours of directed fishing effort. Anglers caught 87 Pumpkinseed and harvested 38. The mean length of harvested fish was 6.7 inches.

ROCK BASS received 3 hours of directed fishing effort. There was no documented catch or harvest of Rock Bass.

ACKNOWLEDGMENTS

The DNR would like to thank all the anglers who took the time to offer information about their fishing trip to the survey clerk. Without their cooperation, the survey would not have been possible.

We also thank our cooperators, Allen and Deanna Hughes, who generously allowed the DNR to keep a boat on their property during this survey.

Completion of this survey was possible because of the efforts of the following fisheries management and treaty fisheries staff: Angelena Sikora, Gene Hatzenbeler, Todd Brecka, Misty Rood, Bill Sobaski, Marty Kangas, Nathan Thomas, Logan Cutler and Samir Shaikh. The creel clerk during the survey period was Chris Coffin.

This creel report was reviewed by Angelena Sikora and Gene Hatzenbeler.

Additional copies of this report, and those covering other local lakes, can be obtained online at:

<http://dnr.wi.gov/topic/Fishing/north/trtycr/srvys.html>

Table 1. Sportfishing effort summary, Amnicon Lake, 2024-25 season; compared to 2006-07 creel results, Douglas County averages, and Ceded Territory averages.

Month	Number of Angler Party Interviews	Total Angler Hours	Total Angler Hours/Acre	2006-07 Total Angler Hours/Acre	Douglas County Average Hours/Acre	Ceded Territory Average Hours/Acre
May	51	672	1.6	3.6	3.6	4.8
June	51	1,165	2.7	4.2	4.2	6.2
July	48	1,343	3.2	3.6	4.4	6.6
August	61	1,220	2.9	3.6	3.1	5.2
September	51	828	1.9	2.3	1.5	3.2
October	21	266	0.6	1.2	0.5	1.4
December	30	656	1.5	1.9	1.3	1.1
January	55	851	2.0	2.5	2.2	1.7
February	57	2,194	5.1	3.1	1.4	1.6
March	8	155	0.4	0.01	0.2	0.2
Summer Total	283	5,495	12.9	18.5	17.4	27.4
Winter Total	150	3,856	9.1	7.5	5.0	4.6
Grand Total	433	9,351	22.0	25.9	22.4	32.0

Note: Summer is May-October; Winter is December-March

Number of Angler Party Interviews is the number of groups of anglers interviewed by the creel clerk. A party is considered the members of a group who fish together in the same boat, ice shanty or from shore. The clerk fills out one interview form for each group of anglers. The number of individual anglers actually contacted by the clerk is usually much greater than the number of groups listed in this table since most groups consist of more than one angler.

Total Angler Hours is the estimated total number of hours that anglers spent fishing on Amnicon Lake during each month surveyed.

Total Angler Hours/Acre is the total angler hours divided by the area of the lake in acres. This is useful in order to compare effort on Amnicon Lake to other lakes.

2006-07 Total Angler Hours/Acre is the total angler hours divided by the area of the lake in acres. This is from the previous creel survey that took place on Amnicon Lake.

County Average Hours/Acre is the average angler effort in hours per acre for county lakes that have been surveyed since 1990. This value is useful for fishing pressure comparisons with other waters.

Ceded Territory Average Hours/Acre is the average angler effort in hours per acre for inland lakes in the Ceded Territory that have been surveyed since 1990. This value can be used to compare Amnicon Lake to other lakes in northern Wisconsin.

Table 2. Comparison of creel survey synopses, Amnicon Lake, 2024-25 and 2006-07 fishing seasons.

CREEL YEAR: 2024-2025

SPECIES	DIRECTED EFFORT (Hours)	PERCENT OF TOTAL	TOTAL CATCH	SPECIFIC CATCH RATE (Hrs/Fish)	TOTAL HARVEST	SPECIFIC HARVEST RATE (Hrs/Fish)	MEAN LENGTH OF HARVESTED FISH
Walleye	928	6.9%	40	35.3	16	59.5	20.9
Northern Pike	4,796	35.6%	4,002	2.0	628	11.2	22.3
Muskellunge	2,162	16.1%	83	27.9	0	*	**
Smallmouth Bass	17	0.1%	0	*	0	*	**
Largemouth Bass	1,679	12.5%	1,012	2.7	35	204.1	15.0
Yellow Perch	38	0.3%	225	*	14	*	6.0
Bluegill	1,273	9.5%	2,925	0.5	564	2.3	6.5
Black Crappie	2,571	19.1%	5,767	0.5	778	3.6	8.0
Pumpkinseed	0	0.0%	87	*	39	*	6.7
Rock Bass	3	0.02%	0	*	0	*	*
Rainbow Smelt	0	0.0%	20	*	0	*	*
Black Bullhead	0	0.0%	3	*	0	*	*

CREEL YEAR: 2006-2007

SPECIES	DIRECTED EFFORT (Hours)	PERCENT OF TOTAL	TOTAL CATCH	SPECIFIC CATCH RATE (Hrs/Fish)	TOTAL HARVEST	SPECIFIC HARVEST RATE (Hrs/Fish)	MEAN LENGTH OF HARVESTED FISH
Walleye	1,958	12.6%	226	45.1	21	495.8	18.0
Northern Pike	166	1.1%	3	1,773.6	0	*	**
Muskellunge	3,917	25.2%	513	17.0	9	963.9	36.0
Smallmouth Bass	180	1.2%	84	65.7	0	*	**
Largemouth Bass	1,832	11.8%	1,300	6.0	8	1,014.4	15.4
Yellow Perch	681	4.4%	2,097	4.3	12	743.1	**
Bluegill	1,844	11.8%	2,951	3.6	54	194.2	6.6
Black Crappie	3,866	24.8%	5,289	2.1	2,264	4.9	8.3
Pumkinseed	1,062	6.8%	1,327	4.8	3	2,049.3	5.3
Rock Bass	63	0.4%	116	43.3	9	555.8	8.5
Shiners	0	0.0%	3	*	0	*	**

Note: If a species is not shown in a table, no data was collected by the creel clerks for that species.

* Indicates that no fish of this species were caught or harvested (depending on the column) by anglers who specifically targeted this species.

** Indicates that no fish were measured by the creel clerks for this species.

WALLEYE









Figure 1. Walleye sportfishing effort, catch, harvest, and length distribution, Amnicon Lake, during 2024-25.



NORTHERN PIKE








Figure 2. Northern Pike sportfishing effort, catch, harvest, and length distribution, Amnicon Lake, during 2024-25.



MUSKELLUNGE







Figure 3. Muskellunge sportfishing effort, catch, harvest, and length distribution, Amnicon Lake, during 2024-25.

SMALLMOUTH BASS








Figure 4. Smallmouth Bass sportfishing effort, catch, harvest, and length distribution, Amnicon Lake, during 2024-25.

LARGEMOUTH BASS












Figure 5. Largemouth Bass sportfishing effort, catch, harvest, and length distribution, Amnicon Lake, during 2024-25.



YELLOW PERCH





Figure 6. Yellow Perch sportfishing effort, catch, harvest, and length distribution, Amnicon Lake, during 2024-25.








Figure 7. Bluegill sportfishing effort, catch, harvest, and length distribution, Amnicon Lake, during 2024-25.

BLACK CRAPPIE









Figure 8. Black Crappie sportfishing effort, catch, harvest, and length distribution, Amnicon Lake, during 2024-25.

PUMPKINSEED









Figure 9. Pumpkinseed sportfishing effort, catch, harvest, and length distribution, Amnicon Lake, during 2024-25.

ROCK BASS








Figure 10. Rock Bass sportfishing effort, catch, harvest, and length distribution, Amnicon Lake, during 2024-25.